

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 April 2004 (08.04.2004)

PCT

(10) International Publication Number
WO 2004/028476 A2

(51) International Patent Classification⁷: **A61K**
(21) International Application Number:
PCT/US2003/030612

18 Kings crossing Court, Apt.J, Baltimore, MD 21030
(US). MCRINER, Andres, J. [GB/US]; 110 W. 39th
Street, Apt.1605, Baltimore, MD 21210 (US).

(22) International Filing Date:
26 September 2003 (26.09.2003)

(74) Agents: PETERSEN, Steven, C. et al.; Hogan & Hart-
son LLP., 1200 17th Street, Suite 1500, Denver, CO 80202
(US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/414,786 27 September 2002 (27.09.2002) US
60/445,526 6 February 2003 (06.02.2003) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): JOHNS
HOPKINS UNIVERSITY [US/US]; Office of Licensing
and Technology Development, 100 Charles Street, 5TH
Floor, Baltimore, MD 21201 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): POSNER, Gary, H.
[US/US]; 3216 Timberfield Lane, Baltimore, MD 21208
(US). SHAPIRO, Theresa, A. [US/US]; 1017 Hart Road,
Towson, MD 21286 (US). SUR, Surojit [IN/US]; 116
W. University Parkway, Apt.1213, Baltimore, MD 21210
(US). LABONTE, Tanzina [BD/US]; 3100 St.Paul Street,
Apt. 415, Baltimore, MD 21218 (US). BORSTNIK,
Kristina [SI/US]; 3811 Canterbury Road, Apt. 711,
Baltimore, MD 21218 (US). PAIK, Ik-Hyeon [KR/US];

Published:

— without international search report and to be republished
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: ORALLY ACTIVE, ANTIMALARIAL, ANTICANCER, ARTEMISININ-DERIVED TRIOXANE DIMERS WITH HIGH SELECTIVITY, STABILITY AND EFFICACY AND METHODS OF MAKING THE SAME

(57) Abstract: In only two steps and in 65% overall yield, natural trioxane artemisinin (I) was converted on gram scale into C-10-carba trioxane dimer (3). This new, very stable dimer was then transformed easily in one additional step into four different dimers (4-7). Alcohol and diol dimers (4 and 5) and ketone dimer (7) are 10 times more antimalarially potent *in vitro* than artemisinin (I), and alcohol and diol dimers (4 and 5) are strongly inhibitory but not cytotoxic toward several human cancer cell lines. Water-soluble carboxylic acid derivatives (8a-10c and 12) were easily prepared from dimers (4-6); they are thermally stable even at 60°C for 24 hours, are more orally efficacious as antimalarials than either artelinic acid or sodium artesunate, and have potent and selective anticancer activities. Further derivitization of the alcohol dimers (4 and 17), diol dimer (5) and ketone (7) has produced a number of analogs also antimalarially active *in vitro* at sub-nanomolar concentrations (most notably: pyridine N-oxides (13, 15, 18, 23, 24 and 25), phosphoric acid triesters (26 and 27), sulfonamide (40) and cyclic carbonate (41)). In addition, dimers (13 and 19) are more efficacious (when administered both orally and i.v.) and less toxic (when administered intraperitoneally to mice as a single dose) than clinically-used sodium artesunate, thereby giving them a better antimalarial therapeutic index than sodium artesunate.